

# Remark on non-Noether symmetries and bidifferential calculi

**George Chavchanidze**

Department of Theoretical Physics, A. Razmadze Institute of Mathematics,  
1 Aleksidze Street, Ge 380093, Tbilisi, Georgia  
e-mail: chaffinch@apexmail.com

**Abstract.** In this paper relationship between non-Noether symmetries and bidifferential calculi is discussed.

In the past few years both non-Noether symmetries [1-4] and bidifferential calculi [5-8] has been successfully used in generating conservation laws and both lead to the similar families of conserved quantities.

Here relationship between Lutzky's integrals of motion [3-4] and bidifferential calculi is briefly discussed.

In the regular Hamiltonian system phase space  $M$  is equipped with symplectic form  $\omega$  (closed  $d\omega = 0$  and non-degenerate  $i_X\omega = 0 \Leftrightarrow X = 0$  2-form) which gives rise to the isomorphism between the vector fields and 1-forms

$$\phi_\omega : TM \longrightarrow T^*M \quad (1)$$

defined by

$$\phi_\omega(X) = i_X\omega \quad (2)$$

where  $i_X\omega$  denotes contraction of the vector field  $X$  with  $\omega$ . This map can be inverted

$$\phi_\omega^{-1} : T^*M \longrightarrow TM \quad (3)$$

with  $\phi_\omega^{-1}(a) = i_W a$   $a \in \Omega^1(M)$  where  $W$  is a bivector field obtained by inverting  $\omega$ . The vector field is said to be Hamiltonian (locally Hamiltonian) if it corresponds to exact (closed) form

$$X_f = \phi_\omega^{-1}(df) \quad (4)$$

According to Liouville's theorem Hamiltonian vector field preserves both  $\omega$  ( $L_{X_f}\omega = 0$  where  $L_{X_f}$  denotes Lie derivative along  $X_f$  vector field) and bivector field  $W$  ( $[X_f, W] = 0$  where  $[ , ]$  is a Schouten bracket).

Now let's consider non-Hamiltonian vector field  $E$  ( $L_E\omega \neq 0$   $[E, W] \neq 0$ ) commuting with the tangent vector of solutions  $[E, X_h] = 0$   $X_h = \phi_\omega^{-1}(dh)$ . The infinitesimal transformation  $g = 1 + \epsilon L_E$  generated by  $E$  is a non-Noether symmetry [3-4] in sense that it doesn't preserve action but maps space of solutions onto itself.

As it was shown by Lutzky [3] such a symmetry leads to the whole family of conserved quantities

$$l_k = i_{W^k} \omega_E^k \quad k = 1, 2, \dots, n \quad (5)$$

(no summation over  $k$ )

where  $\omega_E = L_E \omega$  and  $n$  is dimension of configuration space.

Indeed

$$\frac{d}{dt} l_k = L_{X_h} l_k = L_{X_h} i_{W^k} \omega_E^k = i_{[X_h, W^k]} \omega_E^k + i_{W^k} L_{X_h} \omega_E^k = 0 \quad (6)$$

since  $[X_h, W] = L_{X_h} \omega = 0$  (according to the Liouville's theorem) and  $L_{X_h} \omega_E = L_{X_h} L_E \omega = L_E L_{X_h} \omega = 0$  (due to  $[X_h, E] = 0$ ).

The whole construction can be extended to the singular dynamical systems and to the more general class of symmetries [4].

Now let's turn to the bidifferential calculi. The possibility of generating conservation laws by means of exterior derivative  $d$  and auxiliary differential operator  $d'$  with properties  $d'^2 = 0$  and  $dd' + d'd = 0$  has been considered by Dimakis and Müller - Hoissen [7-8].

According to the Frölicher-Nijenhuis theory action of  $d'$  on functions should be of the following form:  $d'f = i_A df$  where  $A \in \Omega^1(M, TM)$  is tangent valued 1-form with vanishing Frölicher-Nijenhuis torsion [5,6].

$$\begin{aligned} T(A)(X, Y) &= [AX, AY] - A([AX, Y] + [X, AY] - A[X, Y]) = 0 \\ T(A) &\in \Omega^2(M, TM) \quad X, Y \in TM \end{aligned} \quad (7)$$

One can show [5] that traces of powers of  $A$

$$\mu_k = Tr(A^k) \quad (8)$$

form a Lenard scheme

$$d\mu_{k+1} = d'\mu_k \quad (9)$$

and as a result  $\{\mu_k\}$  are in involution - have vanishing Poisson brackets  $\{\mu_k, \mu_{k'}\} = 0$  [6]. This sequence of functions in involution associated with bidifferential calculi  $d, d'$  can be used in generating conservation laws.

Now let's expose relationship between the family of functions  $\{\mu_k\}$  mentioned above and conserved quantities associated with non-Noether symmetry.

By means of the generator of symmetry  $E$  one can construct differential operator  $\hat{d}$  with all properties of  $d'$  besides  $\hat{d}^2 = 0$ . The action of  $\hat{d}$  on functions can be defined by

$$\hat{d}f = \phi_{\omega_E} \circ \phi_{\omega}^{-1}(df) = \hat{A}(f) \quad (10)$$

where  $\hat{A} = (\omega_E)_{ik} W^{kj} dx_i \otimes \frac{\partial}{\partial x_j}$ . Since  $W$  and  $\omega_E$  are invariants ( $L_{X_h} \omega = L_{X_h} W = 0$  according to Liouville's theorem and  $L_{X_h} \omega_E = L_{X_h} L_E \omega = L_E L_{X_h} \omega = 0$ ) therefore operator

$$\phi_{\omega_E} \circ \phi_{\omega}^{-1} : T^*M \longrightarrow T^*M \quad (11)$$

is also constant along solutions and as a result the eigenvalues of  $\phi_{\omega_E} \circ \phi_{\omega}^{-1}$  are integrals of motion.

So if  $\{\lambda_k\}$  are eigenvalues of  $\phi_{\omega_E} \circ \phi_{\omega}^{-1}$  it is easy to verify that Lutzky's conserved quantities  $\{l_k\}$  are expressed by means of  $\{\lambda_k\}$

$$l_k = \sum_{i_s \neq i_{s'}} \lambda_{i_1} \lambda_{i_2} \dots \lambda_{i_k} \quad (12)$$

and vice versa. Traces of powers of  $\hat{A}$  are combinations of  $\{\lambda_k\}$  too

$$\hat{\mu}_k = Tr(\hat{A}^k) = \sum_i \lambda_i^k \quad (13)$$

and therefore are also integrals of motion associated with non-Noether symmetry generated by  $E$  ( $\{\hat{\mu}_k\}$  can be expressed by means of  $\{l_k\}$  and vice versa).

In general we have no restriction on torsion of  $\hat{A}$  (this reflects the fact that in general Lutzky's conserved quantities are not in involution) but when it vanishes  $\{\hat{\mu}_k\}$  reduces to the sequence of functions  $\{\mu_k\}$

$$d\mu_{k+1} = \hat{d}\mu_k \quad (14)$$

associated with bidifferential calculi  $d, \hat{d}$  and considered by Dimakis and Müller-Hoissen.

## Acknowledgments

Author is indebted to Z. Giunashvili for many useful discussions and G. Jorjadze for invaluable help. This work was supported by Scholarship from World Federation of Scientists.

## References

- [1] S. Hojman, A new conservation law constructed without using either Lagrangians or Hamiltonians, 1992 *J. Phys. A: Math. Gen.* 25 L291-295
- [2] F. González-Gascón, Geometric foundations of a new conservation law discovered by Hojman, 1994 *J. Phys. A: Math. Gen.* 27 L59-60
- [3] M. Lutzky, New derivation of a conserved quantity for Lagrangian systems, 1998 *J. Phys. A: Math. Gen.* 15 L721-722
- [4] G. Chavchanidze, Non-Noether symmetries in singular dynamical systems, 2001 *Georgian Math. J.* 8 027-032, math-ph/0106010
- [5] M. Crampin, W. Sarlet, G. Thompson, Bi-differential calculi and bi-Hamiltonian systems, 2000 *J. Phys. A: Math. Gen.* 33 No. 22 L177-180
- [6] P. Guha, A Note on Bidifferential Calculi and Bihamiltonian systems, 2000 *IHÉS preprint* M/64
- [7] A. Dimakis, F. Müller-Hoissen, Bi-differential calculi and integrable models, 2000 *J. Phys. A: Math. Gen.* 33 210-23
- [8] A. Dimakis, F. Müller-Hoissen, Bicomplexes and integrable models, 2000 nlin.SI/0006029
- [9] G. Chavchanidze, Bi-Hamiltonian structure as a shadow of non-Noether symmetry, math-ph/0106018